

Multipolar correlations and deformation effect on nuclear transition matrix elements of double- β decay

R. CHANDRA^{1,2}, K. CHATURVEDI³, P. K. RATH², P. K. RAINA¹ and J. G. HIRSCH⁴

¹ Department of Physics and Meteorology, IIT Kharagpur, Kharagpur-721302, India

² Department of Physics, University of Lucknow, Lucknow-226007, India

³ Department of Physics, Bundelkhand University, Jhansi-284128, India

⁴ Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, A.P. 70-543, México 04510 D.F., México

PACS 21.60.Jz – Hartree-Fock and random-phase approximations

PACS 23.20.-g – Electromagnetic transitions

PACS 23.40.Hc – Relation with nuclear matrix elements and nuclear structure

Abstract. – The two neutrino and neutrinoless double beta decay of $^{94,96}\text{Zr}$, $^{98,100}\text{Mo}$, ^{104}Ru , ^{110}Pd , $^{128,130}\text{Te}$ and ^{150}Nd isotopes for the $0^+ \rightarrow 0^+$ transition is studied within the PHFB framework along with an effective two-body interaction consisting of pairing, quadrupole-quadrupole and hexadecapole-hexadecapole correlations. It is found that the effect of hexadecapolar correlations can be assimilated substantially as a renormalization of the quadrupole-quadrupole interaction. The effect of deformation on nuclear transition matrix elements is investigated by varying the strength of quadrupolar correlations in the parent and daughter nuclei independently. The variation of the nuclear transition matrix elements as a function of the difference in deformation parameters of parent and daughter nuclei reveals that in general, the former tend to be maximum for equal deformation and they decrease as the difference in deformation parameters increases, exhibiting a very similar trend for the $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ transition matrix elements.

The nuclear $\beta\beta$ decay is a rare second order semileptonic transition between two even Z -even N isobars $^A_Z X$ and $^A_{Z\pm 2} Y$ involving strangeness conserving weak charged currents. It is expected that the nuclear $\beta\beta$ decay can proceed through sixteen experimentally distinguishable modes, namely double-electron emission $(\beta^-\beta^-)$, double-positron emission $(\beta^+\beta^+)$, electron-positron conversion $(\varepsilon\beta^+)$ and double-electron capture $(\varepsilon\varepsilon)$ with the emission of two neutrinos, no neutrinos, single Majoron and double Majorons. The $\beta^+\beta^+$, $\varepsilon\beta^+$ and $\varepsilon\varepsilon$ processes are energetically competing and we shall refer to them as $e^+\beta\beta$ modes. The $(\beta\beta)_{2\nu}$ decay conserves the lepton number and is an allowed process in the standard model of electroweak unification (SM). On the other hand, the $(\beta\beta)_{0\nu}$ decay violates the conservation of lepton number and can occur in a number of gauge theoretical models, namely GUTs-left-right symmetric models and $E(6)$ -, R_p -conserving as well as violating SUSY models, in the scenarios of leptoquark exchange, existence of heavy sterile neutrino, compositeness and Majoron models. Hence, it is a convenient tool to test the physics beyond the SM.

The occurrence of $(\beta^-\beta^-)_{2\nu}$ decay was proposed by Mayer in 1935 [1] as the possibility to explain the long observed half-lives of some even-even nuclei and it was finally confirmed by Elliott *et al.* [2] using a TPC detector. Presently, the $(\beta^-\beta^-)_{2\nu}$ decay has been already observed experimentally in ten nuclei, out of 35 possible candidates [3]. For the $(e^+\beta\beta)_{2\nu}$ modes, experimental limits on half-lives have been also given for 24 out of 34 possible isotopes [3]. Klapdor-Kleingrothaus and his group have reported the observation of $(\beta^-\beta^-)_{0\nu}$ decay of ^{76}Ge with $T_{1/2}^{0\nu} = 2.23^{+0.44}_{-0.31} \times 10^{25}$ yr [4, 5]. The first results [4] were debated [6–9]. These results could be confirmed by the forthcoming MAJORANA [10] and GERDA [11] experiments. It would be very exciting if this very first observation of $(\beta^-\beta^-)_{0\nu}$ decay by Klapdor-Kleingrothaus and his coworkers is verified by the latter two mentioned experiments.

It is possible to extract accurate nuclear transition matrix elements (NTMEs) $M_{2\nu}$ from the observed half-lives of $(\beta^-\beta^-)_{2\nu}$ decay for the $0^+ \rightarrow 0^+$ transition and a comparison between the theoretically calculated and experi-

mentally extracted NTMEs provides a cross-check on the reliability of different nuclear models used for the study of nuclear $\beta\beta$ decay. Using the average half-lives [12] and the phase space factors [13], the extracted NTMEs $M_{2\nu}$ vary from 0.0152 ± 0.0008 (0.0238 ± 0.0013) to 0.1222 ± 0.0034 (0.1909 ± 0.0053) for $g_A = 1.25$ (1.00) corresponding to ^{130}Te and ^{100}Mo respectively. In all cases, it is observed that the NTMEs are sufficiently quenched. Vogel and Zirnbauer were the first to provide an explanation of the observed suppression of $M_{2\nu}$ in the QRPA employing a schematic surface-delta interaction including the particle-particle channel, whose strength controls both the concentration of β^- strength in the GTR and the reduction of total β^+ strength and its concentration at low excitation energies [14]. These conclusions were verified by Civitarese *et al.* [15] in the same QRPA model using realistic interaction derived by G-matrix method. In the case of $(\beta^-\beta^-)_{0\nu}$ decay, the nuclear models usually predict half-lives assuming certain value for the neutrino mass, or conversely extract various parameters from the observed limits on half-lives. Over the past years, many a nuclear models have been employed to study the nuclear $\beta\beta$ decay [16,17]. The experimental as well as theoretical study of nuclear $\beta\beta$ decay is quite wide in scope and has been excellently reviewed over the past decades, which can be found in the recent review [18] and references there in.

The structure of a nucleus is mainly decided by the subtle interplay of pairing and quadrupolar correlations present in the effective two-body interaction. The former is responsible for the sphericity of the nucleus, whereas the latter makes the nucleus deformed. In addition to the pairing interaction, which plays an important role in all even Z -even N $\beta\beta$ emitters, it has been also shown that the deformation degrees of freedom are crucial in the structure of ^{100}Mo and ^{150}Nd isotopes [19]. The shell model is the best choice to calculate the NTMEs as it attempts to solve the nuclear many-body problem as exactly as possible. On the other hand, the QRPA and its extensions have emerged as the most successful models in correlating single- β GT strengths and half-lives of $(\beta^-\beta^-)_{0\nu}$ decay. In spite of the spectacular success of the QRPA in the study of $\beta\beta$ decay, there was a need to include the deformation degrees of freedom in its formalism.

Over the past years, the deformed QRPA model has been developed for studying $\beta\beta$ decay of spherical as well as deformed nuclei. The comparison of the experimental GT strength distribution $B(GT)$ with the results of QRPA calculations was employed as a novel method of deducing the deformation of ^{76}Sr nucleus [21]. The effect of deformation on the $(\beta^-\beta^-)_{2\nu}$ decay for the ground state transition $^{76}\text{Ge} \rightarrow ^{76}\text{Se}$ was studied in the framework of the deformed QRPA with separable GT residual interaction [22]. A deformed QRPA formalism to describe simultaneously the energy distributions of the single- β GT strength and the $(\beta^-\beta^-)_{2\nu}$ decay matrix elements of ^{48}Ca , ^{76}Ge , ^{82}Se , ^{96}Zr , ^{100}Mo , ^{116}Cd , $^{128,130}\text{Te}$, ^{136}Xe and ^{150}Nd isotopes using deformed Woods-Saxon poten-

tials and deformed Skyrme Hartree-Fock mean fields was developed [23]. Rodin and Faessler [24] have studied the $\beta\beta$ decay of ^{76}Ge , ^{100}Mo and ^{130}Te isotopes and it has been reported that the effect of continuum on the NTMEs of $(\beta^-\beta^-)_{2\nu}$ decay is negligible whereas the NTMEs of $(\beta^-\beta^-)_{0\nu}$ decay are regularly suppressed.

Recently, large scale calculations in the “Interacting Shell Model” (ISM) have been performed to study the dependence of NTMEs for $\beta^-\beta^-$ decay of ^{48}Ca , ^{76}Ge , ^{82}Se , ^{110}Pd , ^{116}Cd , ^{124}Sn , $^{128,130}\text{Te}$ and ^{136}Xe nuclei on the valence space, effective two-body interactions and deformation [25]. In the ISM, the effect of pairing and quadrupolar correlations on the NTMEs of $(\beta^-\beta^-)_{0\nu}$ decay has been also studied, in which it has been noticed that the NTMEs are the largest in the presence of former only [26,27]. It has been shown by Menendez *et al.* [27] that as the difference of deformation between parent and grand daughter grows, the NTMEs for both $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ decay decrease rapidly.

The PHFB model in conjunction with pairing plus quadrupole-quadrupole (PQQ) interaction [28] is a convenient choice to study the role of quadrupolar correlations vis-a-vis deformation on the NTMEs for $\beta\beta$ decay. However, the PHFB model is unable to provide information about the structure of the intermediate odd-odd nuclei in its present version and hence, on the single β decay rates and the distribution of GT strength. In spite of this limitation, the PHFB model in conjunction with PQQ interaction has been successfully applied to study the $(\beta^-\beta^-)_{2\nu}$ as well as $(e^+\beta\beta)_{2\nu}$ decay modes for the $0^+ \rightarrow 0^+$ transition where it was possible to describe the lowest excited states of the parent and daughter nuclei along with their electromagnetic transition strengths, as well as to reproduce their measured $\beta\beta$ decay rates [29–31].

An inverse correlation between the GT strength and quadrupole moment has been noticed [20]. In the PHFB model, the existence of an inverse correlation between the quadrupole deformation and the size of NTMEs $M_{2\nu}$ has been also confirmed [29–31]. In addition, it has been observed that the NTMEs are usually large in the absence of quadrupolar correlations. With the inclusion of the quadrupolar correlations, the NTMEs are almost constant for small admixture of the QQ interaction and suppressed substantially in realistic situation. In the mass range $A = 90 - 150$, the suppression factor D_α for NTMEs evaluated without and with the quadrupole-quadrupole interaction ranges between 2 and 6, both for the $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ decays. For the double beta decay of ^{150}Nd , it was also found [35] that the NTMEs for $(\beta^-\beta^-)_{0\nu}$ decay have a well defined maximum when the deformation of parent and daughter nuclei are similar and they are suppressed for a difference in the deformation.

The purpose of the present work is two fold. In the first part, we investigate the effect of hexadecapolar correlations on the previously calculated spectroscopic properties, NTMEs $M_{2\nu}$ for $(\beta^-\beta^-)_{2\nu}$ [29,31] and $M_{0\nu}$ for $(\beta^-\beta^-)_{0\nu}$ decay [35]. In the second part, we investigate

the effect of having different deformations in parent and daughter nuclei on NTMEs $M_{2\nu}$ and $M_{0\nu}$ of $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ decay respectively for the $0^+ \rightarrow 0^+$ transition.

The inverse half-life of the $(\beta^-\beta^-)_{2\nu}$ decay for the $0^+ \rightarrow 0^+$ transition is given by [32–34]

$$[T_{1/2}^{2\nu}(0^+ \rightarrow 0^+)]^{-1} = G_{2\nu} |M_{2\nu}|^2 \quad (1)$$

The phase space factor $G_{2\nu}$ can be calculated with good accuracy [13,33] and the nuclear model dependent NTME $M_{2\nu}$ is written as

$$\begin{aligned} M_{2\nu} &= \sum_N \frac{\langle 0_F^+ | \sigma\tau^+ | 1_N^+ \rangle \langle 1_N^+ | \sigma\tau^+ | 0_I^+ \rangle}{E_N - (E_I + E_F)/2} \\ &= \sum_N \frac{\langle 0_F^+ | \sigma\tau^+ | 1_N^+ \rangle \langle 1_N^+ | \sigma\tau^+ | 0_I^+ \rangle}{E_0 + E_N - E_I} \end{aligned} \quad (2)$$

with

$$E_0 = \frac{1}{2} (E_I - E_F) = \frac{1}{2} Q_{\beta\beta} + m_e \quad (3)$$

The expressions to calculate $M_{2\nu}$ in the PHFB model are extensively discussed in our earlier works [29,31,35]. In what follows, we include only the required extensions in the formalism. The HFB wave functions are generated using an effective Hamiltonian with $PQQHH$ type of effective two-body interaction. Explicitly, the Hamiltonian can be written as

$$H = H_{sp} + V(P) + \zeta_{qq} [V(QQ) + V(HH)] \quad (4)$$

where H_{sp} denotes the single particle Hamiltonian. The $V(P)$, $V(QQ)$ and $V(HH)$ represent the pairing, quadrupole-quadrupole and hexadecapole-hexadecapole part of the effective two-body interaction. The ζ_{qq} is an arbitrary parameter and the final results are obtained by setting the $\zeta_{qq} = 1$. The purpose of introducing ζ_{qq} is to study the role of deformation by varying the strength of $QQHH$ interaction.

The pairing part of the effective two-body interaction $V(P)$ is given by

$$V(P) = - \left(\frac{G}{4} \right) \sum_{\alpha\beta} (-1)^{j_\alpha + j_\beta - m_\alpha - m_\beta} a_\alpha^\dagger a_\alpha^\dagger a_\beta a_\beta \quad (5)$$

where α denotes the quantum numbers ($nljm$) and the state $\bar{\alpha}$ is same as α but with the sign of m reversed. The QQ part of the effective interaction $V(QQ)$ is expressed as

$$V(QQ) = - \left(\frac{\chi_2}{2} \right) \sum_{\alpha\beta\gamma\delta} \sum_{\mu} (-1)^\mu \langle \alpha | q_{2\mu} | \gamma \rangle \langle \beta | q_{2-\mu} | \delta \rangle a_\alpha^\dagger a_\alpha^\dagger a_\beta a_\gamma \quad (6)$$

where

$$q_{2\mu} = \left(\frac{16\pi}{5} \right)^{1/2} r^2 Y_{2\mu}(\theta, \phi) \quad (7)$$

The HH part of the effective interaction $V(HH)$ is given as

$$V(HH) = - \left(\frac{\chi_4}{2} \right) \sum_{\alpha\beta\gamma\delta} \sum_{\nu} (-1)^\nu \langle \alpha | q_{4\nu} | \gamma \rangle \langle \beta | q_{4-\nu} | \delta \rangle a_\alpha^\dagger a_\alpha^\dagger a_\beta a_\gamma \quad (8)$$

with

$$q_{4\nu} = r^4 Y_{4\nu}(\theta, \phi) \quad (9)$$

The relative magnitudes of the parameters of the HH part of the two body interaction are calculated from a relation suggested by Bohr and Mottelson [39]. According to them the approximate magnitude of these constants for isospin $T = 0$ is given by

$$\chi_\lambda = \frac{4\pi}{2\lambda + 1} \frac{m\omega_0^2}{A \langle r^{2\lambda-2} \rangle} \quad \text{for } \lambda = 1, 2, 3, 4 \dots \quad (10)$$

and the parameters for the $T = 1$ case are approximately half of their $T = 0$ counterparts. The value of χ_4 for $T = 1$ is taken as exactly half of the $T = 0$ case. Explicitly

$$\begin{aligned} \chi_4 &= \left[\left(\frac{16}{25} \right) \left(\frac{2}{3} \right)^{2/3} \right] \chi_2 A^{-2/3} b^{-4} \\ &= 0.4884 \chi_2 A^{-2/3} b^{-4} \end{aligned} \quad (11)$$

with $b = 1.0032 A^{1/6}$.

The model space, single particle energies (SPE's), parameters of PQQ type of effective two-body interactions and the method to fix them are the same as in references [29,31,35]. We fix χ_{2pn} as well as χ_{4pn} through the experimentally available energy spectra for a given model space, SPE's, G_p , G_n , χ_{2pp} and χ_{4pp} . The strength of proton-neutron (pn) component of the QQ interaction χ_{2pn} is varied so as to obtain the spectra of considered nuclei, namely $^{94,96}\text{Zr}$, $^{94,96,98,100}\text{Mo}$, $^{98,100,104}\text{Ru}$, $^{104,110}\text{Pd}$, ^{110}Cd , $^{128,130}\text{Te}$, $^{128,130}\text{Xe}$, ^{150}Nd and ^{150}Sm in optimum agreement with the experimental results. The theoretical spectra has been taken to be the optimum one if the excitation energy of the 2^+ state E_{2^+} is reproduced as closely as possible to the experimental value. All these input parameters are kept fixed to calculate other spectroscopic properties.

In table 1, we present the values of χ_{2pn} for the PQQ and $PQQHH$ interactions. It is clear from the fitted values of χ_{2pn} that once the procedure to fix E_{2^+} is adopted, the change in the latter is minimal even in the presence of HH correlations. This behavior is further reflected in the calculated yrast spectra, reduced $B(E2:0^+ \rightarrow 2^+)$ transition probabilities, static quadrupole moments $Q(2^+)$ and gyromagnetic factors $g(2^+)$, which are usually in an overall agreement with the experimental data [29,31]. The maximum change in all the calculated spectroscopic properties is about 10% except for the $B(E2:0^+ \rightarrow 2^+)$ and $Q(2^+)$ of ^{94}Zr , which change by 20% and 23% respectively. Further, we tabulate the theoretically calculated and experimental deformation parameter β_2 [36] of parent and daughter nuclei involved in $\beta^-\beta^-$ decay in the same table 1 at $\zeta_{qq} = 1$. The calculated NTMEs $M_{2\nu}$ for the $0^+ \rightarrow 0^+$ transition at $\zeta_{qq} = 1$ are given in table 2. The change in results for β_2 are below 10%. The NTMEs $M_{2\nu}$ also change up to 10% except ^{94}Zr , ^{130}Te and ^{150}Nd , for which the changes are 17%, 26% and 21% respectively. The ratio for deformation effect $D_{2\nu}$ are 2.29(2.23), 3.70(3.38),

Table 1: Fitted values of χ_{2pn} , calculated deformation parameters β_2 of parent(P) and daughter(D) nuclei participating in $\beta^-\beta^-$ decay using PQQ and $PQQHH$ type of two-body interaction along with the average experimental values. † and ‡ denote PQQ and $PQQHH$ interaction respectively.

		χ_{2pn}		β_2	
			Theory	Experiment [36]	
^{94}Zr	†	0.02519	0.100	0.090 ± 0.010	
	‡	0.02629	0.110		
^{94}Mo	†	0.02670	0.161	0.1509 ± 0.0015	
	‡	0.02572	0.161		
^{96}Zr	†	0.01717	0.085	0.080 ± 0.017	
	‡	0.01918	0.087		
^{96}Mo	†	0.02557	0.191	0.1720 ± 0.0016	
	‡	0.02472	0.186		
^{98}Mo	†	0.01955	0.158	0.1683 ± 0.0028	
	‡	0.01955	0.162		
^{98}Ru	†	0.02763	0.205	0.1947 ± 0.0030	
	‡	0.02649	0.194		
^{100}Mo	†	0.01906	0.231	0.2309 ± 0.0022	
	‡	0.01876	0.226		
^{100}Ru	†	0.01838	0.214	0.2148 ± 0.0011	
	‡	0.01831	0.214		
^{104}Ru	†	0.02110	0.285	0.2707 ± 0.0020	
	‡	0.02053	0.282		
^{104}Pd	†	0.01486	0.216	0.209 ± 0.007	
	‡	0.01507	0.219		
^{110}Pd	†	0.01417	0.216	0.257 ± 0.006	
	‡	0.01393	0.214		
^{110}Cd	†	0.01412	0.196	0.1770 ± 0.0039	
	‡	0.01414	0.191		
^{128}Te	†	0.02715	0.136	0.1363 ± 0.0011	
	‡	0.02692	0.136		
^{128}Xe	†	0.03600	0.192	0.1836 ± 0.0049	
	‡	0.02662	0.181		
^{130}Te	†	0.01801	0.117	0.1184 ± 0.0014	
	‡	0.01890	0.120		
^{130}Xe	†	0.02454	0.166	0.169 ± 0.007	
	‡	0.02281	0.163		
^{150}Nd	†	0.02160	0.276	0.2853 ± 0.0021	
	‡	0.02228	0.279		
^{150}Sm	†	0.01745	0.238	0.1931 ± 0.0021	
	‡	0.01730	0.241		

1.86(1.75), 2.33(2.09), 5.47(5.19), 3.14(3.30), 4.26(3.97), 2.89(3.31) and 5.94(6.46) for $^{94,96}\text{Zr}$, $^{98,100}\text{Mo}$, ^{104}Ru , ^{110}Pd , $^{128,130}\text{Te}$ and ^{150}Nd nuclei with $PQQ(PQQHH)$ interaction respectively and their variation due to the $PQQHH$ type of interaction is up to 15%.

A suppression of NTMEs $M_{2\nu}$ for $\beta\beta$ decay with respect to the spherical case has been reported when the parent and daughter nuclei have different deformations [23, 27]. To investigate this effect, we present the NTMEs for the $(\beta^-\beta^-)_{2\nu}$ as well as $(\beta^-\beta^-)_{0\nu}$ decay of $^{94,96}\text{Zr}$, $^{98,100}\text{Mo}$, ^{104}Ru , ^{110}Pd isotopes in fig. 1 and $^{128,130}\text{Te}$ and ^{150}Nd

Table 2: Calculated NTMEs $M_{2\nu}$ and $M_{0\nu}$ for $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ decay respectively using PQQ [29, 31, 35] and $PQQHH$ interaction along with the extracted $M_{2\nu}$ from average/recommended experimental half-lives for the $0^+ \rightarrow 0^+$ transition [12], except for ^{94}Zr [37], and ^{110}Pd [38]. The numbers corresponding to (a) and (b) are calculated for $g_A = 1.25$ and 1.0 respectively. † and ‡ denote PQQ and $PQQHH$ interaction respectively.

		$M_{2\nu}$		$ M_{0\nu} $
		Theory	Experiment	Theory
			(a)	(b)
^{94}Zr	†	0.076	<62.8	<98.1
	‡	0.063		
^{96}Zr	†	0.058	0.051	0.080
	‡	0.059		
^{98}Mo	†	0.130		
	‡	0.127		
^{100}Mo	†	0.104	0.122	0.191
	‡	0.104		
^{104}Ru	†	0.068		
	‡	0.068		
^{110}Pd	†	0.133	<6.47	<10.11
	‡	0.120		
^{128}Te	†	0.033	0.022	0.034
	‡	0.033		
^{130}Te	†	0.042	0.015	0.024
	‡	0.031		
^{150}Nd	†	0.033	0.033	0.051
	‡	0.026		

isotopes in fig. 2 as a function of the difference in the deformation parameter $\Delta\beta_2 = \beta_2(\text{parent}) - \beta_2(\text{daughter})$ between the parent and daughter nuclei. The NTMEs are calculated by keeping the deformation for parent nuclei fixed at $\zeta_{qq} = 1$ and changing the deformation of daughter nuclei by varying ζ_{qq} in the range 0.0 – 1.5.¹

In our earlier works [29, 31], it has been shown that the NTMEs $M_{2\nu}$ are usually large for $\zeta_{qq} = 0.0$ i.e. when both the parent and daughter nuclei are spherical. With the increase of ζ_{qq} , the NTMEs remain almost constant and then decrease around the physical value $\zeta_{qq} = 1.0$ establishing an inverse correlation between $M_{2\nu}$ and β_2 . It can be observed at the left panels of figs. 1 and 2 that in all cases that when the absolute value of the difference in deformation is close to zero the largest NTMEs are obtained. In addition, the effect of hexadecapolar correlations is almost negligible.

In the mass mechanism neglecting pseudo-scalar and weak magnetism terms of the recoil current, the importance of which has been reported by Simkovic *et al.* [40, 41] and needs further investigation, the inverse half-life of the

¹In table 5 of ref. [31], the given β_2 values of ^{130}Xe (for $^{130}\text{Te} \rightarrow ^{130}\text{Xe}$) at $\zeta_{qq} = 0.9$ and 1.5 are 0.159 and 2.155 respectively, which is a typographical error. The correct values are 0.157 and 0.216 respectively.




Fig. 1: NTMEs of $^{94,96}\text{Zr}$, $^{98,100}\text{Mo}$, ^{104}Ru , ^{110}Pd isotopes for $(\beta^-\beta^-)_{2\nu}$ (left hand side) and $(\beta^-\beta^-)_{0\nu}$ (right hand side) decay as a function of the difference in the deformation parameter $\Delta\beta_2$. “x” denotes the value of NTME for calculated $\Delta\beta_2$ from table 1.

$(\beta^-\beta^-)_{0\nu}$ decay due to the exchange of light neutrinos for the $0^+ \rightarrow 0^+$ transition is given by [32–34]

$$\left[T_{1/2}^{0\nu}(0^+ \rightarrow 0^+)\right]^{-1} = \left(\frac{\langle m_\nu \rangle}{m_e}\right)^2 G_{01} |M_{0\nu}|^2 \quad (12)$$




Fig. 2: NTMEs of $^{128,130}\text{Te}$ and ^{150}Nd nuclei for $(\beta^-\beta^-)_{2\nu}$ and $(\beta^-\beta^-)_{0\nu}$ decay. Further details are given in fig. 1.

where the NTME $M_{0\nu}$ is given by

$$M_{0\nu} = \sum_{n,m} \left\langle 0_F^+ \left\| \left[\sigma_1 \cdot \sigma_2 - \left(\frac{g_V}{g_A} \right)^2 \right] H_m(r) \tau_n^+ \tau_m^+ \right\| 0_I^+ \right\rangle \quad (13)$$

The calculation of $M_{0\nu}$ in the PHFB model has been discussed in an earlier work [35]. It is worth mentioning that we include the finite size and short range effects through a dipole form factor and Jastrow type of short range correlation (SRC). The other two prescriptions for including SRC namely, through the exchange of ρ and ω mesons [42] and unitary correlation operator method [43, 44] need to be investigated. In table 2, we give the theoretically calculated $M_{0\nu}$ using the HFB wave functions in conjunction with PQQ [35] and $PQQHH$ interaction. The change in the values of NTMEs $M_{0\nu}$ are below 10% except ^{110}Pd , ^{130}Te and ^{150}Nd , for which the changes are 17%, 18% and 25% respectively. The ratio for deformation effect $D_{0\nu}$ are 2.51(2.30), 4.50(4.21), 1.94(1.87), 2.15(1.92), 3.81(3.85), 2.64(2.87), 4.43(3.85), 2.96(3.26) and 6.16(7.14) for $^{94,96}\text{Zr}$, $^{98,100}\text{Mo}$, ^{104}Ru , ^{110}Pd , $^{128,130}\text{Te}$ and ^{150}Nd nuclei with $PQQ(PQQHH)$ interaction respectively corresponding to a maximum change up to 16%. It is worth mentioning that, the β_2 for ^{150}Nd is close to the experimental value whereas for ^{150}Sm , it is off. For the PQQ interaction, the experimental β_2 but for other physical properties can be reproduced with $\zeta_{qq} = 0.96$ and the corresponding $M_{0\nu}$ is 0.7015.

In the right panel of figs. 1 and 2, we present the variation of $|M_{0\nu}|$ with respect to $|\Delta\beta_2|$ for the above mentioned $\beta^-\beta^-$ emitters. It is noticed that variation in $|M_{0\nu}|$ with changing $|\Delta\beta_2|$ is similar to that of $(\beta^-\beta^-)_{2\nu}$ decay.

Moreover, it is noticed from figs. 1 and 2 that the NTMEs remain constant even when one of the nuclei is spherical or slightly deformed. With further increase in deformation, the NTMEs become the maximum for $|\Delta\beta_2| = 0$ and then decrease with increase in the difference between the deformation parameter.

We conclude that the effect of including the hexadecapole interaction channel is minor both on the deformation β_2 and on the NTMEs. In particular, it is clear that the qualitative dependence of the NTMEs on deformation is not affected by including the HH interactions in the Hamiltonian, and the changes, while visible, are small. It exhibits that the inclusion of QQ interaction is a necessary ingredient for a reliable calculation of NTMEs, and that effect of the hexadecapolar correlations can be mimicked by a slight renormalization in the QQ channel. The independent deformations of initial and final nuclei are, indeed, crucial parameters to describe realistic NTMEs $M_{2\nu}$ and $M_{0\nu}$ of $\beta\beta$ decay, which exhibit, in their respective scales, a very similar dependence on the difference of deformations.

* * *

Partial grant by DST, India (SR/S2/HEP-13/2006), Conacyt-México and DGAPA-UNAM is acknowledged.

REFERENCES

- [1] GOEPFERT-MAYER M., *Phys. Rev.*, **48** (1935) 512.
- [2] ELLIOTT S. R., HANN A. A. and MOE M. K., *Phys. Rev. Lett.*, **59** (1987) 2020; **59** (1987) 1649.
- [3] TRETYAK V. I. and ZDESENKO Y. G., *At. Data Nucl. Data Tables*, **61** (1995) 43; **80** (2002) 83.
- [4] KLAPDOR-KLEINGROTHAUS H. V. *et al.*, *Mod. Phys. Lett. A*, **16** (2001) 2409.
- [5] KLAPDOR-KLEINGROTHAUS H. V., KRIVOSHEINA I., DIETZ A. and CHKVORETS O., *Phys. Lett. B*, **586** (2004) 198; KLAPDOR-KLEINGROTHAUS H. V. and KRIVOSHEINA I., *Mod. Phys. Lett. A*, **21** (2006) 1547.
- [6] FERUGLIO F., STRUMIA A. and VISSANI F., *Nucl. Phys. B*, **637** (2002) 345.
- [7] AALSETH C. E. *et al.*, *Mod. Phys. Lett. A*, **17** (2002) 1475.
- [8] ZDESENKO YU. G., DANEVICH F. A. and TRETYAK V. I., *Phys. Lett. B*, **546** (2002) 206.
- [9] IANNI A., *Nucl. Instrum. Meth. A*, **516** (2004) 184.
- [10] GEHMAN V. M. *et al.*, *AIP Conf. Proc.*, **842** (2006) 840.
- [11] BETTINI A. and THE GERDA COLLABORATION, *Nucl. Phys. B (Proc. Suppl.)*, **168** (2007) 67.
- [12] BARABASH A. S., *Czech. J. Phys.*, **56**, (2006) 437.
- [13] BOEHM F. and VOGEL P., *Ann. Rev. Nucl. Part. Sci.*, **34** (1984) 125; *Physics of Massive Neutrinos, 2nd ed.* (Cambridge University Press, Cambridge) 1992.
- [14] VOGEL P. and ZIRNBAUER M. R., *Phys. Rev. Lett.*, **57** (1986) 3148.
- [15] CIVITARESE O., FAESSLER A. and TOMODA T., *Phys. Lett. B*, **194** (1987) 11.
- [16] SUHONEN J. and CIVITARESE O., *Phys. Rep.*, **300** (1998) 123.
- [17] FAESSLER A. and SIMKOVIC F., *J. Phys. G*, **24** (1998) 2139.
- [18] AVIGNONE F. T., ELLIOTT S. R. and ENGEL J., *Rev. Mod. Phys.*, **80** (2008) 481.
- [19] GRIFFITHS A. and VOGEL P., *Phys. Rev. C*, **46** (1992) 181; SUHONEN J. and CIVITARESE O., *Phys. Rev. C*, **49** (1994) 3055.
- [20] AUERBACH N., ZHENG D. C., ZAMICK L. and BROWN B. A., *Phys. Lett. B*, **304** (1993) 17; AUERBACH N., BERTSCH G. F., BROWN B. A. and ZHAO L., *Nucl. Phys. A*, **556** (1993) 190; TROLTENIER D., DRAAYER J. P. and HIRSCH J. G., *Nucl. Phys. A*, **601** (1996) 89.
- [21] NÁCHER E. *et al.*, *Phys. Rev. Lett.*, **92** (2004) 232501.
- [22] PACEARESCU L., FAESSLER A. and SIMKOVIC F., *Phys. At. Nucl.*, **67** (2004) 1210.
- [23] ÁLVAREZ-RODRÍGUEZ R., SARRIGUREN P., MOYA DE GUERRA E., PACEARESCU L., FAESSLER A. and SIMKOVIC F., *Phys. Rev. C*, **70** (2004) 064309; ÁLVAREZ-RODRÍGUEZ R., SARRIGUREN P., MOYA DE GUERRA E., PACEARESCU L., SAAKYAN R. and SIMKOVIC F., *Prog. Part. Nucl. Phys.*, **57** (2006) 251.
- [24] RODIN V. and FAESSLER A., *Phys. Rev. C*, **77** (2008) 025502.
- [25] CAURIER E., NOWACKI F. and POVES A., *Eur. Phys. J. A*, **36** (2008) 195.
- [26] CAURIER E., MENENDEZ J., NOWACKI F. and POVES A., *Phys. Rev. Lett.*, **100** (2008) 052503.
- [27] MENENDEZ J., POVES A., CAURIER E. and NOWACKI F., *arxiv:0809.2183 [nucl-th]*, (2008) .
- [28] BARANGER M. and KUMAR K., *Nucl. Phys. A*, **110** (1968) 490.
- [29] CHANDRA R., SINGH J., RATH P. K., RAINA P. K. and HIRSCH J. G., *Eur. Phys. J. A*, **23** (2005) 223.
- [30] RAINA P. K., SHUKLA A., SINGH S., RATH P. K. and HIRSCH J. G., *Eur. Phys. J. A*, **28** (2006) 27.
- [31] SINGH S., CHANDRA R., RATH P. K., RAINA P. K. and HIRSCH J. G., *Eur. Phys. J. A*, **33** (2007) 375.
- [32] HAXTON W. C. and STEPHENSON G. J., JR., *Prog. Part. Nucl. Phys.*, **12** (1984) 409.
- [33] DOI M., KOTANI T. and TAKASUGI E., *Prog. Theor. Phys.*, **83** (1985) 1.
- [34] TOMODA T., *Rep. Prog. Phys.*, **54** (1991) 53.
- [35] CHATURVEDI K., CHANDRA R., RATH P. K., RAINA P. K. and HIRSCH J. G., *Phys. Rev. C*, **78** (2008) 054302.
- [36] RAMAN S., NESTOR C. W. and TIKKANEN JR. P., *At. Data Nucl. Data Tables*, **78** (2001) 1.
- [37] ARNOLD R. *et al.*, *Nucl. Phys. A*, **658** (1999) 299.
- [38] WINTER R. G., *Phys. Rev.*, **85** (1952) 687.
- [39] BOHR A. and MOTTELSON B. R., *Nuclear Structure*, Vol. **1** (World Scientific, Singapore) 1998.
- [40] SIMKOVIC F., PANTIS G., VERGADOS J. D., and FAESSLER A., *Phys. Rev. C*, **60** (1999) 055502.
- [41] VERGADOS J. D., *Phys. Rep.*, **133** (1986) 1; **361** (2002) 1.
- [42] HIRSCH J. G., CASTAÑOS O. and HESS P. O., *Nucl. Phys. A*, **582** (1995) 124.
- [43] KORTALINEN M. and SUHONEN J., *Phys. Rev. C*, **76** (2007) 024315; KORTALINEN M., CIVITARESE O., SUHONEN J. and TOIVANEN J., *Phys. Lett. B*, **647** (2007) 128.
- [44] SIMKOVIC F., FAESSLER A., RODIN V., VOGEL P. and ENGEL J., *Phys. Rev. C*, **77** (2008) 045503.